Pesticide Safety Fact Sheet



ONIONS

Pesticides commonly used on **onions** in Nevada are listed below. Pesticides may be applied any time during the growing season. If you mix, load, or apply pesticides, or if you work in areas where pesticides have been applied you need to know

how to protect yourself from pesticide illness. Do not enter fields that have posted warning signs, and never allow children to enter treated areas. The chart below tells the number of hours (restricted entry interval) that you should stay out of a field after a pesticide was applied. Always read the pesticide label for information about restricted entry and field posting.

ed gn	Si Pe Po
	•
	•
	•
	•
	•
	•
	•
	•
	Kee _l skin
	slee



Keep out of the field if you see this sign.

1 = Most Dangerous 4 = Least Dangerous	Field Posted Warning Sign
1. DANGER! POISON! PELIGRO! Dyfonate (48 hrs) Lannate (48 hrs)	Yes No
2. DANGER! PELIGRO! Warrior (24 hrs)	No
3. WARNING! AVISO! Lorsban 4E (24 hrs) Telone II (5 days) Ridomil (24 hrs) Goal (24 hrs) Bravo (48 hrs) Ambush (12 hrs) Buctril (12 hrs)	No Yes No No No No
4. CAUTION! Fusilade (12 hrs) Rovral (24 hrs) Benlate (24 hrs) Prowl (24 hrs) Dithane (24hrs)	No No No No No

Signs of Pesticide Poisoning

rashes or itching

watery eyes

blurry vision

headaches

dizziness

chest pain

sweating

vomiting

Keep pesticides off your skin by wearing long sleeved shirts, long pants, shoes, socks and protective gloves.
Pesticides stay on your clothes. Wash your work clothes every time you wear them. Clean work clothes in a different load from other laundry. Wash your hands with soap before eating, drinking, smoking, chewing gum or tobacco or going to the bathroom.

If you or someone else gets sick while working-

- tell your boss right away
- the phone number of a medical facility is listed on the safety poster at your work
- the sick person should be driven to the hospital (don't let them drive them self)
- your boss needs to give the name of the pesticide to the doctor